
Citation:

Nicholson, B and Jones, B and Dinsdale, A and Till, K (2019) Relationships between jump and sprint force-velocity profiles and performance. In: 2019 NSCA National Conference, 10 July 2019 - 13 July 2019, Washington DC.

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/6222/>

Document Version:

Conference or Workshop Item (Presentation)

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

RELATIONSHIPS BETWEEN JUMP AND SPRINT FORCE-VELOCITY PROFILES AND PERFORMANCE

Ben Nicholson^{1,3}, Ben Jones^{1,2,3,4,5,6}, Alex Dinsdale¹ & Kevin Till^{1,2,3}



@Nicholsonsandc



B.T.Nicholson@leedsbeckett.ac.uk

Purpose

To investigate the associations between matched mechanical variables derived from both vertical¹ and horizontal² force-velocity-power (FVP) profiling, and the performance outcome variables within squat jump (SJ) and sprint performance.

Method



20 elite male academy rugby league players (age 17.6 ± 0.9 years; height 179.9 ± 6.6 cm; body mass 91.2 ± 11.8 kg).

- Data collection:** The participants performed two maximal 40m sprints. Sprints were recorded using a radar gun device (Stalker ATS II), which obtained instantaneous speed-time measurements. Sprint times were determined from the modelled velocity-time data at 2m, 5m, 10m and 20m and maximum velocity (V_{max} ; $m \cdot s^{-1}$)².
- The participants performed two maximal SJ (~90° knee angle) repetitions with 0kg, 20kg, 40kg, 60kg and 80kg. An Optojump was used to record jump height (cm) for each load.
- Body mass relative vertical and horizontal mechanical variables (theoretical maximal values of force (F_0) (N/kg), velocity (V_0) (m/s), power (P_{max}) (W/kg)) and the slope of the F-V linear relationship (S_{fv}) were calculated^{1,2}.
- Data analysis:** Pearson's correlation coefficients (r) assessed the relationship between matched vertical and horizontal mechanical variables (F_0 vertical & horizontal, v_0 vertical & horizontal, P_{max} vertical & horizontal and S_{fv} vertical & horizontal) and SJ and sprint performance.



Figure 1&2. Mechanical profiling methods imagery

Results

Table 1. Horizontal and vertical mechanical variables Mean \pm SD data.

	F_0 (N/kg)	V_0 (m/s)	S_{fv} (N.s/m/kg)	P_{max} (W/kg)
HZT variable	7.31 ± 1.04	8.7 ± 0.54	-0.84 ± 0.13	15.9 ± 2.58
VTC variable	31.34 ± 5.15	4.16 ± 1.51	-8.8 ± 4.42	31.51 ± 9.6

Table 2. Performance variable Mean \pm SD data.

	Jump height (cm)	2m time (s)	5m time (s)	10m time (s)	20m time (s)	V_{max} (m/s)
Performance variable	32.34 ± 11.59	0.82 ± 0.06	1.41 ± 0.08	2.17 ± 0.12	3.49 ± 0.18	8.42 ± 0.49

Table 3. Pearson correlation between matched mechanical variables

	F_0 (N/kg)	V_0 (m/s)	S_{fv} (N.s/m/kg)	P_{max} (W/kg)
HZT & VTC	$r = -0.19$	$r = 0.15$	$r = -0.25$	$r = 0.34$

Table 4. Pearson correlation between HZT mechanical and performance variables.

	HZT- F_0 (N/kg)	HZT- V_0 (m/s)	HZT- S_{fv} (N.s/m/kg)	HZT- P_{max} (W/kg)
Jump height (cm)	$r = -0.42$	$r = -0.25$	$r = -0.36$	$r = -0.43$
2m time (s)	$r = -0.98^{***}$	$r = -0.22$	$r = -0.83^{***}$	$r = -0.94^{***}$
5m time (s)	$r = -0.96^{***}$	$r = -0.32^*$	$r = -0.78^{***}$	$r = -0.96^{***}$
10m time (s)	$r = -0.93^{***}$	$r = -0.42^*$	$r = -0.71^{***}$	$r = -0.98^{***}$
20m time (s)	$r = -0.85^{***}$	$r = -0.59^{***}$	$r = -0.59^{***}$	$r = -0.98^{***}$
V_{max} (m/s)	$r = 0.17$	$r = 1.0^{***}$	$r = 0.26$	$r = 0.55^{***}$

Table 5. Pearson correlation between VTC mechanical and performance variables.

	VTC- F_0 (N/kg)	VTC- V_0 (m/s)	VTC- S_{fv} (N.s/m/kg)	VTC- P_{max} (W/kg)
Jump height (cm)	$r = -0.13$	$r = 0.47^*$	$r = 0.41$	$r = 0.51^*$
2m time (s)	$r = 0.16$	$r = -0.40$	$r = -0.27$	$r = -0.39$
5m time (s)	$r = 0.19$	$r = -0.40$	$r = -0.28$	$r = -0.37$
10m time (s)	$r = 0.17$	$r = -0.35$	$r = -0.25$	$r = -0.32$
20m time (s)	$r = 0.21$	$r = -0.39$	$r = -0.30$	$r = -0.36$
V_{max} (m/s)	$r = -0.05$	$r = 0.17$	$r = 0.12$	$r = 0.16$

Note: F_0 , theoretical maximal force; v_0 , theoretical maximal velocity; P_{max} , theoretical maximal power; HZT, horizontal; VTC, vertical. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

Results Summary

- There was no significant correlation between vertical and horizontal FVP matched mechanical variables ($p > 0.05$).
- The correlations between vertical FVP variables and sprint performance and between horizontal FVP variables and SJ performance failed to reach statistical significance ($p > 0.05$).
- Moderate -0.32 to near perfect 1.0 significant correlations ($p < 0.05$) were found between mechanical and performance variables shifting the importance of separate variables depending on the testing task.




Figure 3 & 4. Example mechanical profiles sprint and SJ³

Conclusions

- The absence of significant correlations between the vertical and horizontal FVP profiles suggests that they provide distinctive information about the athlete's mechanical variables.
- The magnitude of the correlations between mechanical variables and sprint performance shifted across the velocity-time curve, therefore performance is determined by separate qualities depending on the distance.
- Whereas, P_{max} reported the greatest correlation with SJ height.

Practical Applications

- To ensure specific, accurate and comprehensive characterisation of athletes' physical qualities FVP profiles should be determined with exercises maximal mechanically similarity to the targeted performance task.
- These results will aid practitioners in test selection the prescription and individualisation of training by providing important information as to the most influential variables to develop SJ and sprint performance.

Acknowledgments

The travel and conference fees were funded by Leeds Beckett University

References

- Samozino, P., Morin, J.B., Hittay, E. and Belli, A., 2008. A simple method for measuring force, velocity and power output during squat jump. *Journal of biomechanics*, 41(14), pp.2940-2945.
- Samozino, P., Rabita, G., Doré, S., Slawinski, J., Peyrot, N., Saez de Villarreal, E. and Morin, J.B., 2016. A simple method for measuring power, force, velocity properties, and mechanical effectiveness in sprint running. *Scandinavian journal of medicine & science in sports*, 26(6), pp.648-658.
- Jiménez-Reyes, P., Samozino, P., García-Ramos, A., Cuadrado-Perfíñez, V., Brughe, M. and Morin, J.B., Relationship between vertical and horizontal force-velocity-power profiles in various sports and levels of practice.